To learn more about your plan, please see anthem.com.

To learn more about vaccines, please see the Centers for Disease Control and Prevention (CDC) website: cdc.gov.

Anthem Blue Cross and Blue Shield is the trade name of: In Colorado: Rocky Mountain Hospital and Medical Service, Inc. HMO products underwritten by HMO Colorado, Inc. In Connecticut: Anthem Health Plans, Inc. In Indiana: Anthem Insurance Companies, Inc. In Kentucky: Anthem Health Plans of Kentucky, Inc. In Maine: Anthem Health Plans of Maine, Inc. In Missouri (excluding 30 counties in the Kansas City area): RightCHOICE® Managed Care, Inc. (RIT), Healthy Alliance® Life Insurance Company (HALIC), and HMO Missouri, Inc. RIT and certain affiliates administer non-HMO benefits underwritten by HALIC and HMO benefits underwritten by HMO Missouri, Inc. RIT and certain affiliates only provide administrative services for self-funded plans and do not underwrite benefits. In Nevada: Rocky Mountain Hospital and Medical Service, Inc. HMO products underwritten by HMO Colorado, Inc., dba HMO Nevada. In New Hampshire: Anthem Health Plans of New Hampshire, Inc. In Ohio: Community Insurance Company. In Virginia: Anthem Health Plans of Virginia, Inc. trades as Anthem Blue Cross and Blue Shield in Virginia, and its service area is all of Virginia except for the City of Fairfax, the Town of Vienna, and the area east of State Route 123. In Wisconsin: Blue Cross Blue Shield of Wisconsin (BCBSWi), which underwrites or administers the PPO and indemnity policies; Compcare Health Services Insurance Corporation (Compcare), which underwrites or administers the HMO policies; and Compcare and BCBSWi collectively, which underwrite or administer the POS policies. Independent licensees of the Blue Cross and Blue Shield Association. ® ANTHEM is a registered trademark of Anthem Insurance Companies, Inc. The Blue Cross and Blue Shield names and symbols are registered marks of the Blue Cross and Blue Shield Association.

Preventive health guidelines

As of April 2011

What is your plan for better health?

Make this year your best year for wellness. Your health plan may help pay for tests to find disease early and routine wellness exams to help you and your family stay well. Talk with your doctor (health care provider) about the care that is right for you.

Your plan may not pay for all services and treatments in this guide. To learn more about what your plan pays for, see your certificate of coverage or call the customer care number on your ID card. You also can check **anthem.com** to learn about health topics from child care to zinc.

The content in this guide is based in part on suggestions from these independent groups:

- · AAFP American Academy of Family Physicians
- · AAP American Academy of Pediatrics
- · ACIP Advisory Committee on Immunization Practices
- · ACOG American College of Obstetrics and Gynecology
- · ACS American Cancer Society
- · ATP III/NCEP Adult Treatment Panel III/National Cholesterol Education Program
- · CDC Centers for Disease Control and Prevention
- USPSTF U.S. Preventive Services Task Force

This guide is just for you to learn from, it is not meant to take the place of medical care or advice.

Use this guide to know when to set up visits with your doctor for you and your children. Ask your doctor which exams, tests and vaccines are right for you, when you should get them and how often. How you get certain diseases is not talked about in this guide.

Please see your plan handbook to check on your plan benefits.



Baby and child screenings

Well-baby exam — birth to 2 years*: Infants who leave the hospital less than two days (48 hours) after birth need to be seen by a doctor within two to four days after being born. Also, you might want to talk to a doctor before your baby is born. You might talk to the doctor if you are a first-time parent, are having a high-risk pregnancy, or want to learn about feeding, circumcision or well-baby care. At the well-baby exam you may get advice on your child's safety, health, healthy eating and development. At these exams, your baby may get vaccines and these screenings:

	Age (in months)									
Screenings	birth	1	2	4	6	9	12	15	18	24
Hearing		as	a new	born a	nd whe	n your	doctor	sugges	sts	
Weight, length and head circumference (the length around the head)	at each visit									
Hemoglobin or hematocrit (blood count)	t) once between 9 and 12									
Lead testing unless you are sure the child has not been around lead							at 12			at 24
Development and behavior	at each visit									
Oral/dental health					at 6	at 9	at 12		at 18	at 24
Autism									at 18	at 24

Well-child exam — ages 2¹/₂ **to 10 years*:** You may get advice about how to keep your child safe, how to prevent injuries, good health, healthy eating and development. At ages 3 and 6, a referral to a dentist may be suggested. At these well-child exams, your child may get vaccines and these screenings:

	Age (in years)								
Screenings	2 ¹ / ₂	3	4	5	6	7	8	9	10
Blood pressure	each year starting at 3 years								
Vision	each year								
Hearing	each year								
Height, weight, body mass index (BMI)**	each year								
Development and behavior	at each visit								
Oral/dental health	at 2 ¹ / ₂								

Well-child exam — ages 11 to 18 years*: The doctor may talk to you about health and wellness issues. These include healthy eating, exercise, healthy weight, how to prevent injuries, avoiding tobacco, alcohol and drugs, sexual behavior, dental health, mental health and secondhand smoke. At these exams, your child may get vaccines and these screenings:

	Age (in years)									
Screenings	11	12	13	14	15	16	17	18		
Blood pressure		each year								
Vision	each year									
Hearing	each year									
Height, weight, BMI**	each year									
Chlamydia	for sexually active women aged 25 and younger									
Development and behavior	havior each year									

^{*}This guide is for people with average risk. Some people may be at higher risk for health issues due to their family health history, their race or ethnicity, or other reasons. Talk to your doctor if you have concerns about your health.

4-6 years	11-12 years	13-18 years	19-49 years	50-64 years	65+ years
V					
	✓ Tdap		Td boos	ter every	10 years
~					
from 6 m	onths to	65+ years	of age		
~					
V					
	3-dose series				
	V				
			sugge for smol asthmat	kers and	1 lifetime dose
				1 singl for age	e dose es 60+

✓ Shows when vaccines are suggested

Tdap (teens) - If you are 13 to 18 years of age and have not had this vaccine before, talk to your doctor about a catch-up vaccine.

Td (adults) - Adults should have one dose of Tdap.

Flu - Refer to flu.gov or cdc.gov to learn more about this vaccine. (Note: Children 6 months to 8 years of age having the vaccine for the first time should have two doses.)

Measles, mumps, rubella (MMR) and varicella - If you are 19 years of age or older and you have no proof you had these vaccines and you have never had these illnesses, then you should get these vaccines.

HPV - This vaccine is for girls 11 to 12 years of age, but it may be given as early as 9 years of age. Girls and women aged 13 through 26 who have not had the vaccine, should talk to their doctor about a catch-up vaccine. HPV may be given to males aged 9 through 26 years to lessen their chance of getting genital warts.

Meningococcal vaccine - Two doses of this vaccine are given. The first dose at 11 or 12 years of age and the next dose at 16.

Pneumococcal conjugate (PCV) - The suggested vaccine has changed. Children 14 to 59 months who had all of the PCV vaccine series, may need an extra vaccine shot at their next visit.*

*Children 14 to 59 months who had all of the PCV vaccine series with 7-valent pneumococcal conjugate vaccine (PCV7) should get 1 more dose of 13-valent pneumococcal conjugate vaccine (PCV13) at their next visit.

^{**}Height and weight is used to find BMI. BMI for children is based on gender and age. It is used to see if a person is underweight, overweight or at risk for overweight. BMI for adults is used to see if a person is overweight or obese.

Suggested vaccine schedule*

							Age		
Vaccine	birth	1-2 months	2 months	4 months	6 months	6-18 months	12-15 months	15-18 months	19-23 months
Hepatitis B	V	/		see note		V			
Rotavirus (RV)			✓ 2-dose	e or 3-dose	series**				
Diphtheria, tetanus, pertussis (DTaP)			V	~	V			~	
Tetanus, diphtheria, pertussis (Td/Tdap)									
Haemophilus influenza type b (Hib)			✓ 3-	-dose or 4-	dose serie	S***			
Pneumococcal conjugate (PCV)			V	~	V		~		
Inactivated polio virus (IPV)			~	~		•			
Influenza							✓ su	ggested e	each year
Measles, mumps, rubella (MMR)							~		
Varicella (chicken pox)							V		
Hepatitis A							V	2-dose ser	ies
Human papillomavirus (HPV)									
Meningococcal									
Pneumococcal polysaccharide (PPSV)									
Zoster									
	Note: Ge	t this vacci	ne at this a	nge, only if t	the combir	nation vacc	ine is used	after the b	irth dose.

^{*}This guide is for people with average risk. Some people may be at higher risk for health issues due to their family health history, their race or ethnicity, or other reasons. Talk to your doctor if you have concerns about your health.

Adult screenings — women*

Well-person exam: The doctor may talk with you about health and wellness issues. These include healthy eating, exercise, family planning for ages 19-39 and folic acid for women who are of the age to get pregnant, how to prevent injuries, misuse of drugs and alcohol, how to stop using tobacco, secondhand smoke, sexual behavior, dental health and mental health. At this visit, you may get vaccines and these screenings:

	Age (in years)										
Screenings	19 20 25 30 35 40 45 50 55 60 65 and older										
Blood pressure	each year or as your doctor suggests										
Height, weight	each year or as your doctor suggests										
BMI	each year or as your doctor suggests										
Cholesterol	every 5 years starting at age 20 with more screenings as your doctor suggests										
Colorectal cancer	at age 50, your doctor may suggest one of these test options: 1. Fecal occult blood test (FOBT) each year 2. Flexible sigmoidoscopy every 5 years 3. Both #1 and #2 4. Double-contrast barium enema every 5 years 5. Colonoscopy every 10 years 6. CT colonography may take the place of a colonoscopy in some cases										
Hearing	when doctor suggests										
Breast cancer: self-exam	each month from age 19 to 65+										
Breast cancer: doctor exam	every 2 to 3 years each year from age 40 to 65+										
Breast cancer: mammogram	each year from age 40 to 65+										
Cervical cancer: ages 19-39	each year starting within 3 years of sexual activity or age 21 (whichever comes first)										
Cervical cancer: ages 30-69	every 2 to 3 years for women with 3 normal Pap tests in a row										
Cervical cancer: ages 70+	with 3 or more normal Pap tests in a row and no abnormal Pap tests in the last 10 years, women may stop having this screening										
Chlamydia	sexually active women ages 25 and younger										
Osteoporosis	the test to check how dense your bones are should start no later than age 65; women at menopause should talk to their doctor about osteoporosis and have the test when at risk										

^{**}Rotavirus - 2-dose or 3-dose series; depends on brand of vaccine used

^{***}Haemophilus influenza type b (Hib) - 3-dose or 4-dose series; depends on brand of vaccine used

^{****}Pneumococcal polysaccharide (PPSV) for smokers and asthmatics – give 1 dose if never got the vaccine or if don't know vaccine history

^{*}This guide is for people with average risk. Some people may be at higher risk for health issues due to their family health history, their race or ethnicity, or other reasons. Talk to your doctor if you have concerns about your health.

Pregnant women*

Pregnant women should see their doctor or OB/GYN in their first three months for a first visit and to set up a prenatal care plan. At this visit your doctor will check your health and the health of your baby.

Based on your past health, your doctor may want you to have these tests, screenings or vaccines:

- · Hematocrit/hemoglobin (blood count)
- · Syphilis
- · HIV
- · Rubella immunity to find out which women need the rubella vaccine after giving birth
- · Rh(D) blood type and antibody testing if Rh(D) negative, repeat test at 26 to 28 weeks
- · Hepatitis B
- · Urinalysis when your doctor wants it

The doctor may talk to you about what to eat and how to be active when pregnant as well as staying away from tobacco, drugs, alcohol and other substances.

Other tests and screenings:

Some tests given alone or with other tests can be used to check the baby for health concerns. These tests are done at certain times while you are pregnant. The best test to use and the best time to do it depends on many things. These include your age as well as your medical and family history. Talk to your doctor about what these tests can tell you about your baby, the risks of the tests and which tests may be best for you.

- · Special blood tests
- **Ultrasound tests** including special tests (used with blood tests during the first three months for chromosomal abnormality risk) and routine two-dimensional tests to check on the baby
- · Amniocentesis
- · Chorionic villus sampling

Vaccines:

If you are pregnant in flu season (October to March) your doctor may want you to have the inactivated flu vaccine. You may need the tetanus, diphtheria, pertussis (Tdap) vaccine if you have not had it before. While other vaccines may be given in special cases, it is best to get the vaccines you need before you get pregnant. Women should always check with their doctor about their own needs.

You should NOT get these vaccines while you are pregnant:

- · Measles, mumps, rubella (MMR)
- · Varicella

Adult screenings — men*

Well-person exam: The doctor may talk with you about health and wellness issues. These include healthy eating, exercise, family planning for ages 19-39, how to prevent injuries, misuse of drugs and alcohol, how to stop using tobacco, secondhand smoke, sexual behavior, dental health and mental health. At this visit, you may get vaccines and these screenings:

							Age (in yea	ırs)			
Screenings	19	20	25	30	35	40	45	50	55	60	65 and older	
Blood pressure		each year or as your doctor suggests										
Height, weight					each	ı year (or as y	our do	ctor su	ıggest	'S	
BMI					each	n year (or as y	our do	ctor su	ıggest	CS .	
Cholesterol		every 5 years starting at age 20 with more screenings as your doctor suggests									re screenings	
Colorectal cancer						at age 50, your doctor may suggest one of these test options: 1. Fecal occult blood test (FOBT) each year 2. Flexible sigmoidoscopy every 5 years 3. Both #1 and #2 4. Double-contrast barium enema every 5 year 5. Colonoscopy every 10 years 6. CT colonography may take the place of a colonoscopy in some cases						
Prostate cancer								if you are 50 or older, discuss with your doctor the risks and benefits of the prostate cancer tests				
Abdominal aortic aneurysm											one time for ages 65-75 for those who have ever smoked	
Hearing											when your doctor suggests	

^{*}This guide is for people with average risk. Some people may be at higher risk for health issues due to their family health history, their race or ethnicity, or other reasons. Talk to your doctor if you have concerns about your health.

^{*}This guide is for people with average risk. Some people may be at higher risk for health issues due to their family health history, their race or ethnicity, or other reasons. Talk to your doctor if you have concerns about your health.